

US006291725B1

(12) United States Patent

Chopade et al.

(10) Patent No.: US 6,291,725 B1

(45) **Date of Patent:** Sep. 18, 2001

(54) CATALYSTS AND PROCESS FOR HYDROGENOLYSIS OF SUGAR ALCOHOLS TO POLYOLS

(75) Inventors: Shubham P. Chopade, East Lansing;
Dennis J. Miller, Okemos; James E.
Jackson, Haslett, all of MI (US); Todd
A. Werpy, West Richland, WA (US);
John G. Frye, Jr.; Alan H. Zacher,
both of Richland, WA (US)

(73) Assignees: Board of Trustees operating Michigan
State University, East Lansing, MI
(US); Battelle Memorial Institute,
Pacific Northwest Laboratory,
Richland, WA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 09/518,860

(22) Filed: Mar. 3, 2000

(51) **Int. Cl.**⁷ **C07C 31/18**; C07C 29/60

(52) **U.S. Cl.** **568/861**; 568/862; 568/863; 568/864

(56) References Cited

U.S. PATENT DOCUMENTS

1,963,997	6/1934	Larchar .
2,004,135	6/1935	Rothrock .
2,852,570	9/1958	Conradin et al.
3,030,429	4/1962	Conradin et al.
3,396,199	8/1968	Kasehagen .
4,338,472	7/1982	Sirkar .
4,366,332	12/1982	Chao et al
4,380,675	4/1983	Gebauer et al.
4,401,823	8/1983	Arena .
4,404,411	9/1983	Tanikella .

4,430,253	2/1984	Dubeck et al
4,496,780	1/1985	Arena .
4,552,862	* 11/1985	Larkin 502/306
5,026,927	6/1991	Andrews et al
5,210,335	5/1993	Schuster et al
5,326,912	7/1994	Gubitosa et al
5,354,914	10/1994	Gubitosa et al
5,600,028	2/1997	Gubitosa et al

OTHER PUBLICATIONS

Van Ling and Vlugter, J. Appl. Chem. 19:43 (1969). Muller et al in Heterogeneous Catalysis and Fine Chemicals II, M. Guisnet et al, eds. Elsevier Press, Amsterdam, Netherlands, p. 237 (1991).

Montassier et al., J. Mole. Catal. 70: 99 (1991). Brunauer, emmett, and Teller (BET), J. Am. Chem. Soc., 60:309 (1938).

* cited by examiner

Primary Examiner—Johann Richter
Assistant Examiner—Brian J. Davis

(74) Attorney, Agent, or Firm—Ian C. McLeod

(57) ABSTRACT

The present invention provides a process for preparation of low molecular weight polyols from high molecular weight polyols in a hydrogenolysis reaction under elevated temperature and hydrogen pressure. The process comprises providing in a reaction mixture the polyols, a base, and a metal catalyst prepared by depositing a transition metal salt on an inert support, reducing the metal salt to the metal with hydrogen, and passivating the metal with oxygen, and wherein the catalyst is reduced with hydrogen prior to the reaction. In particular, the process provides for the preparation of glycerol, propylene glycol, and ethylene glycol from sugar alcohols such as sorbitol or xylitol. In a preferred process, the metal catalyst comprises ruthenium which is deposited on an alumina, titania, or carbon support, and the dispersion of the ruthenium on the support increases during the hydrogenolysis reaction.

23 Claims, 3 Drawing Sheets